

# Particle-Size Analysis

Geochemistry of the Land - Dr. C.E. Heinzl  
University of Northern Iowa  
Dept. of Earth and Environmental Sciences

# Learning objectives

2

- Review sedimentary particle textures
- Know Udden-Wentworth grain size scale and Phi ( $\phi$ ) scale, and the relationships between the two
- Know how to plot grain size data: histograms; frequency curves; cumulative frequency curves
- Know how to obtain grain size data
- Understand grain size descriptive statistics
  - Mode
  - Median
  - Graphic estimation of mean
  - Phi standard deviation (graphic estimation of sorting)
  - Graphic skewness

# Sedimentary textures

3

Textural properties of siliciclastic sediments and rocks include:

1. Grain size
2. Sorting
3. Grain shape
  1. Sphericity
  2. Rounding
4. Particle surface textures
5. Grain fabric

# Grain size scales

## Udden-Wentworth scale

- Geometric scale in which each step is twice as large as the previous one
- Range is from 1/256 mm to >256 mm
- Four major size classes are
  - Gravel (> 2.00 mm)
  - Sand (1/16 mm to 2.00 mm)
  - Silt (1/256 mm to 1/16 mm)
  - Clay (< 1/256 mm)

} mud

# Grain size scales (continued)

5

- Phi scale (Krumbein)

- Logarithmic scale useful for plotting and statistical calculations


$$\Phi = -\log_2 d$$

(where  $d$  = grain diameter, in mm)

Note that 1.00 mm = 0 $\phi$

$\Phi$  increases with smaller grain size,  
decreases with larger grain size

# Handout




Wentworth grain size chart from United States Geological Survey Open-File Report 2006-1195

# Measuring grain size

7

- Technique for measuring grain size depends on:
  - Purpose of study
  - Size range of material being measured
  - Consolidated vs. unconsolidated
- Grain size is usually expressed in terms of long dimension or intermediate dimension of particle



# Measuring grain size (continued)

8

Type of sample	Sample grade	Method of analysis
Unconsolidated sediment and disaggregated sedimentary rock	Boulders Cobbles Pebbles	Manual measurement of individual clasts
	Granules Sand Silt	Sieving, settling-tube analysis, image analysis
	Clay	Pipette analysis, sedimentation balances, photohydrometer, Sedigraph, laser-diffractometer, electro-resistance (e.g., Coulter counter)
Lithified sedimentary rock	Boulders Cobbles Pebbles	Manual measurement of individual clasts
	Granules Sand Silt	Thin-section measurement, image analysis
	Clay	Electron microscope



# Measuring grain size (continued)

9

Sieving is useful for unconsolidated material ranging from granule to silt size

- Sieve screen catches particles according to intermediate dimension

Petrographic microscope with ocular micrometer is useful for lithified sand and silt sized material

- Actual measurements or visual comparison with known standards
- Measurements tend to underestimate actual size because grain axes rarely lie in plane of thin section

# Graphical treatment of grain size data

10

Common graphical methods for presenting grain size data are:

- Histogram

- Individual weight percent is plotted for each  $\phi$  size class

- Frequency curve

- A histogram in which a smooth curve connects the midpoints of each size class

- Cumulative curve

- A plot of  $\phi$  grain size versus cumulative weight percent
- Most useful for estimating statistical parameters


# Graphical treatment of grain size data

11

$\phi$ size class	raw weight (gm)	individual weight percent	cumulative weight percent
-1.00	0.39	0.50	0.50
-0.50	1.96	2.50	3.00
0.00	3.92	5.00	8.00
0.50	6.26	8.00	16.00
1.00	8.61	11.00	27.00
1.50	11.75	15.00	42.00
2.00	12.53	16.00	58.00
2.50	11.75	15.00	73.00
3.00	8.61	11.00	84.00
3.50	6.26	8.00	92.00
4.00	3.92	5.00	97.00
4.50	1.96	2.50	99.50
5.00	0.39	0.50	<b>100.00</b>
<b>total</b>	<b>78.3</b>		


# Histogram

12




# Frequency curve

13



# Cumulative curve



Steep curve indicates good sorting; flat curve indicates poor sorting

# Grain size statistics

15

**Mode** = most frequently occurring particle size in a population of grains

- Highest point of a frequency curve or histogram; steepest segment of a cumulative curve

**Median** = midpoint of grain size distribution

- 50<sup>th</sup> percentile on cumulative curve

**Mean** = arithmetic average grain size


- Practically impossible to determine
- Can be approximated graphically

**Mean = median = mode** in a perfectly normal distribution; **but**, most natural samples are not normally distributed!

# Determining the graphic mean particle size:

16

$$M_Z = \frac{[\phi_{16} + \phi_{50} + \phi_{84}]}{3}$$





# Sorting


17

Sorting is a measure of the range of grain sizes present in a population and the magnitude of the scatter of those sizes around the mean size


Sorting can be estimated by visual comparison with known standards or by graphic calculations

# Visual estimation of sorting


18




VERY WELL SORTED



WELL SORTED



MODERATELY SORTED





POORLY SORTED

# Graphic estimation of sorting

$$\sigma_i = \frac{\phi_{84} - \phi_{16}}{4} + \frac{\phi_{95} - \phi_5}{6.6}$$

## Phi standard deviation

- < 0.35 $\phi$  = very well sorted
- 0.35-0.50 $\phi$  = well sorted
- 0.50-0.71 $\phi$  = moderately well sorted
- 0.71-1.00 $\phi$  = moderately sorted
- 1.00-2.00 $\phi$  = poorly sorted
- 2.00-4.00 $\phi$  = very poorly sorted
- > 4.00 $\phi$  = extremely poorly sorted



# Application and importance of grain size data

20

Grain size is most useful as a descriptive property insofar as it relates to the derived properties


- porosity and permeability

Grain size, in and of itself, is not a reliable indicator of depositional environments because the grain size distribution of a sediment is the result of processes not environments, but it may infer the energy present

- i.e., sediment transport processes are not unique to any given environment

# Skewness

measure of the amount of departure from a normal distribution (i.e., an asymmetrical frequency curve)



# Graphical estimation of skewness

22

## Graphic skewness

$$SK_i = \frac{\phi_{84} + \phi_{16} - 2\phi_{50}}{2(\phi_{84} - \phi_{16})} + \frac{\phi_{95} + \phi_5 - 2\phi_{50}}{2(\phi_{95} - \phi_5)}$$

>0.30

strongly fine skewed

0.30 to 0.10

fine skewed

0.10 to -0.10

nearly symmetrical

-0.10 to -0.30

coarse skewed

< -0.30

strongly coarse skewed

# Stoke's Law

23

$$V = \frac{(\rho_s - \rho_f)gR^2}{18\mu}$$

V = terminal velocity

$\rho_s$  = Mass density of particles ( $\text{kg}/\text{m}^3$ )

$\rho_f$  = Mass density of fluid ( $\text{kg}/\text{m}^3$ )

$\mu$  = Viscosity of water ( $\text{kg}/\text{m}^*\text{s}$ )

g = gravitational acceleration ( $\text{m}/\text{s}^2$ )

D = diameter of sphere

- Note that if the particle and the fluid have identical densities, no settling will occur.
- Stoke's Law is valid only for particles smaller than 0.2mm

# Comparing and contrasting methods

- Pipetting and sieving
- Sieving



# AVG. Results

24

## Wentworth Geometric Progression

## USDA

Sample	Sand (mm)						Silt (mm)						Clay	Sand	Silt	Clay		
	VCS	CS	MS	FS	VFS	TS	VCSi a	VCSi b	VCSi	CSi	MSi	FSi	VFSi	TSi	TC	TS	TSi	TC
1	11.0	7.3	3.7	2.9	3.3	28.1	0.7	3.5	4.2	14.7	5.7	5.8	6.7	37.0	34.9	28.8	36.3	34.9
2	3.7	9.4	14.2	13.3	8.8	49.4	1.4	3.1	4.5	11.9	2.2	2.6	2.8	24.1	26.5	50.8	22.7	26.5
3	19.0	25.7	22.2	11.7	5.0	83.6	0.4	0.8	1.2	7.4	-0.2	1.7	1.1	11.1	5.2	84.0	10.7	5.2

1 - Black sediment (C)

2 - Brown sediment (B)


3 - Red sediment (D)

Data						raw weights(g)	individual weight %	cumulative Weight %	raw weights		
						A1	A2				
						56.8	32.45	32.45	63.52		
						23.25	13.28	45.74	32.51		
						49.91	28.52	74.25	27.98		
						38.61	22.06	96.31	30.44		
						3.99	2.28	98.59	11.75		
						0.16	0.09	98.68	3.27		
						0.31	0.18	98.86	5.19		
						175.03			174.66		

Phi units (Φ)	A1	A2	A3	A4	A5	mean	median	mode	st deviation
-2.0	33.59	36.37	3.38	0.00	2.06	15.08	3.38	3.38	16.31
-1.0	46.88	54.98	22.89	0.06	5.54	26.07	22.89	22.89	21.80
0.0	45.39	71.00	41.82	24.87	26.82	41.98	41.82	41.82	16.59
1.0	95.45	88.43	59.6	97.84	84.70	85.204	88.43	88.43	13.64
2.0	99.73	95.16	63.3	99.93	98.75	91.374	98.75	98.75	14.14
3.0	99.82	97.03	65.03	99.97	99.53	92.276	99.53	99.53	13.67
4.0	100.00	100.00	100	100.00	100.00	100	100.00	100.00	


# Basic histograms

Sample A1





entire data set



# cumulative weights

